

Neutralino Relic Density in the CPVMSSM at the ILC

G. Bélanger¹, O. Kittel², S. Kraml³, H. U. Martyn⁴ and A. Pukhov⁵

1- LAPTH, Univ. de Savoie, CNRS
B.P. 110, F-74941 Annecy-le-vieux - France

2- Departamento de Física Teórica y del Cosmos and CAFPE,
Universidad de Granada, E-18071 Granada, Spain

3- LPSC, Univ. Joseph-Fourier, CNRS/IN2P3, INPG
53 avenue des Martyrs, F-38026 Grenoble - France

4- I. Physikalisches Institut, RWTH Aachen
Sommerfeldstrasse 14, D-52074 Aachen - Germany

5- SINP, Moscow State Univ.
Moscow 119992 - Russia

We discuss ILC measurements for a specific MSSM scenario with CP phases, where the lightest neutralino, a candidate for dark matter, annihilates through t-channel exchange of light staus. These prospective ILC measurements are used to fit the underlying model parameters. A collider prediction of the relic density of the neutralino from this fit gives $0.116 < \Omega h^2 < 0.19$ at 95% CL.

1 Introduction

One of the prime motivations for a high-luminosity e^+e^- linear collider is the possibility to do precision measurements of new particles beyond the Standard Model and in particular of the dark matter (DM) candidate. A precise determination of the properties of the DM in the laboratory could be used to make a “collider prediction” of its relic abundance, which can be tested against cosmological models. For such a collider prediction to be of interest, it must be at least as precise as the value obtained from cosmological observations, about 10% for WMAP+SDSS [2] and a few percent at the PLANCK satellite [3].

The possibility to make collider predictions of the cross sections for annihilation of dark matter candidates has been examined within specific supersymmetric scenarios. For example, in the general MSSM, it was shown that in a favourable scenario the LHC could match roughly the WMAP+SDSS precision [4] while the ILC could achieve much better precision [5]. These conclusions, however, depend very strongly on the scenario considered; many remain challenging even at the ILC [5].

These studies assumed that CP is conserved, although CP-violating (CPV) phases are generic in the MSSM. CPV MSSM phases can have an important effect on the neutralino annihilation cross sections [6] and could lead to variations in Ωh^2 solely from modifications in the couplings of up to an order of magnitude [7]. Therefore the determination of the relevant couplings (including phases) can be as important for the prediction of Ωh^2 as measurements of masses. In a particular scenario of the CPV MSSM, the precision to which the underlying MSSM parameters could be determined from measurements at the ILC was investigated in [8]. The resulting precision on the neutralino relic density that could be inferred from this was examined. The main results are summarised below.

2 The stau-bulk scenario of the CPVMSSM

The scenario we investigate belongs to the “stau bulk” region of [7], which appears for light staus and large phase of M_1 , the gaugino mass. The neutralino LSP is dominantly bino and annihilates predominantly into tau pairs, via t-channel exchange of both $\tilde{\tau}_1$ and $\tilde{\tau}_2$. The annihilation cross section is sensitive to the stau mixing. Although the $\tilde{\tau}_1$ is very light, the scenario does not rely on coannihilation (the mass difference with the LSP is too large).

As a benchmark point we choose the following input parameters at the electroweak scale:

$$\begin{aligned} M_1 &= 80.47 \text{ GeV}, & M_2 &= 170.35 \text{ GeV}, & M_3 &= 700 \text{ GeV}, & \phi_1 &= \pi, & \mu &= 600 \text{ GeV} \\ M_{\tilde{L}_3} &= 138.7 \text{ GeV}, & M_{\tilde{E}_3} &= 135.2 \text{ GeV}, & A_\tau &= 60 \text{ GeV}, & \phi_\tau &= 0, & \tan \beta &= 10. \end{aligned} \quad (1)$$

Other sfermion masses, $A_{t,b}$, and $m_{H^\pm}$ are set to 1 TeV and $\phi_\mu = 0$. This way EDM constraints are avoided when varying ϕ_1 and ϕ_τ . The particles accessible at ILC500 are $\tilde{\chi}_1^0$, $\tilde{\chi}_2^0$, $\tilde{\chi}_1^\pm$, $\tilde{\tau}_1$, $\tilde{\tau}_2$ and $\tilde{\nu}_\tau$ with masses respectively of (80.7, 164.9, 164.9, 100.9, 177.2, 123.1) GeV as computed with CPsuperH [9]. Despite having a light spectrum this scenario is quite challenging for colliders. Production of sparticles at ILC500 only lead to τ ’s plus E_T^{miss} and similarly for the cascade decays of squarks at LHC.

The relic density of the $\tilde{\chi}_1^0$ computed with micrOMEGAs2.2 [10] is $\Omega h^2 = 0.130$. The precision with which Ωh^2 can be inferred from ILC measurements depends not only on the accuracy of the sparticle spectroscopy but also on the determination of all parameters of the neutralino sector ($M_1, M_2, \mu, \tan \beta, \phi_1$) and of the stau sector ($M_{\tilde{\tau}_L}, M_{\tilde{\tau}_R}, A_\tau, \phi_\tau$). The dependence on ϕ_1 , and to a much lesser extent ϕ_τ , originates from the $\tilde{\chi}_1^0 \tilde{\tau}_{1,2} \tau$ couplings. In addition, particles which are too heavy to be produced at ILC could have some influence. We assume here that the mass scale of the squarks and gluino is known from LHC and that the ILC excludes selectrons/smuons and heavy Higgs states up to the kinematic limit. This leaves an overall uncertainty from the unknown part of the spectrum of $\delta\Omega/\Omega \simeq 7\%$.

3 ILC measurements

channel	observables
$\tilde{\tau}_1^+ \tilde{\tau}_1^-$	$m_{\tilde{\tau}_1} = 100.92 \pm 0.40 \text{ GeV}$ $m_{\tilde{\chi}_1^0} = 80.67 \pm 0.35 \text{ GeV}$ $\cos 2\theta_{\tilde{\tau}} = -0.065 \pm 0.028$ $\mathcal{P}_\tau = 0.64 \pm 0.035$
$\tilde{\tau}_2^+ \tilde{\tau}_2^-$	$m_{\tilde{\tau}_2} = 176.9 \pm 9.1 \text{ GeV}$
$\tilde{\chi}_1^+ \tilde{\chi}_1^-$	$m_{\tilde{\chi}_1^\pm} = 164.88 \pm 0.015 \text{ GeV}$

Table 1: Achievable precisions at ILC.

$\cos 2\theta_{\tilde{\tau}}$, can be determined from measurements of the polarized cross section while the measurement of the τ polarisation $\mathcal{P}_\tau$ in the decay $\tilde{\tau}_1 \rightarrow \tau \tilde{\chi}_1^0$ gives additional information on the stau and neutralino mixings [11]. It is in particular useful to constrain the gaugino–higgsino composition of the LSP. It was shown in [8] that masses, polarisation $\mathcal{P}_\tau$ and mixing $\theta_{\tilde{\tau}}$ can be accurately determined even with moderate integrated luminosity, see Table 1. This analysis used simulations based on SIMDET 4.02 [12], beam polarisations $(\mathcal{P}_{e^-}, \mathcal{P}_{e^+}) = (0.8, 0.6)$ and includes QED radiation, beamstrahlung and SM background to W pairs.

At the ILC, sparticle masses can be measured either with threshold scans or with the endpoint method. The challenge in this scenario is to disentangle the various sources that lead to the $\tau\tau E_T^{\text{miss}}$ topology. Scanning downwards in energy in steps of 10 GeV while using different beam polarisations allows to detect thresholds: the σ_{RL} polarisation mode for $\tilde{\tau}_i^+ \tilde{\tau}_j^-$ pair production and the σ_{LR} mode for $\tilde{\chi}^+ \tilde{\chi}^-$ production. In addition the mixing angle in the stau sector,

4 DM properties: fit to ILC observables

We performed a fit to the ILC observables listed in Table 1 and used a Markov Chain Monte Carlo (MCMC) method to probe the 8-dimensional parameter space. The free parameters are M_1 , μ , $\tan\beta$, $M_{\tilde{L}_3}$, $M_{\tilde{R}_3}$, A_τ , ϕ_1 , ϕ_τ , while M_2 is computed from μ and $\tan\beta$ to match the extremely well measured value of $m_{\tilde{\chi}_1^\pm}$. The total $\tau\tau$ SUSY cross section with polarized beams was also included in the fit. Moreover, we have $\sigma_{\tau\tau} = 3.220 \pm 0.046$ pb at $\sqrt{s} = 400$ GeV including the systematic and statistical uncertainties for 10 fb^{-1} .

The very precise determination of $m_{\tilde{\chi}_1^0}$ and $m_{\tilde{\chi}_1^\pm}$ constrain M_1 to $\delta M_1 = -0.9, +2.1$ GeV and indicates a correlation between ϕ_1 and M_1 although the phase ϕ_1 can take any value. The measurement of the stau masses and their mixing angle constrain the stau soft masses to about 10 GeV. The trilinear coupling A_τ and its phase ϕ_τ are on the other hand basically undetermined. This is because the stau mixing which is proportional to $(A_\tau - \mu \tan\beta)$ is dominated by the term $\mu \tan\beta$. This also means that μ and $\tan\beta$ are individually poorly determined, see Fig. 1.

Within the allowed parameter space, the relic density of dark matter is predicted to be $0.116 < \Omega h^2 < 0.19$ at 2σ . The largest values of Ωh^2 are found for $\phi_1 \approx 0$ and small μ , where the LSP has an increased higgsino component, see Fig. 2. An improvement on the limit on the neutralino–proton cross section, which is largest when μ is small, would cut on these scenarios thus somewhat reducing the range for Ωh^2 .

A better determination of $\tan\beta$ and μ would be needed to significantly improve the




Figure 1: 2σ (green) and 1σ (black) allowed region in the $\tan\beta$ vs μ plane.




Figure 2: Predictions for Ωh^2 vs a) μ and b) ϕ_1 and the WMAP upper limit (line).

uncertainty in Ωh^2 . For both, a multi-TeV e^+e^- linear collider offers the best prospects: The parameter μ could be determined from a measurement of the heavy higgsino states. A measurement of the stop masses and mixing angle at LHC and a multi-TeV LC would completely determine the stop sector thus reducing the large parametric uncertainty in m_h . The precise determination of m_h at the ILC could then be used to constrain $\tan\beta$.

CP-odd observables such as electric dipole moments or T-odd asymmetries could give a very clear signal of CP violation. However, it was shown in [8] that a T-odd asymmetry based on triple-products in the production and decay of neutralinos [13] while sensitive to ϕ_1 , would not help constrain Ωh^2 . The reason is a two-fold ambiguity in ϕ_1 .

5 Conclusions

A precise determination of SUSY particle properties in the “stau-bulk” scenario of the CPVMSSM is quite challenging at the ILC. Although some of the underlying Lagrangian parameters can be extracted with very good precision, the large uncertainties in μ , $\tan\beta$ and the phase ϕ_1 induce a large uncertainty in the prediction of the neutralino relic density. To reduce this uncertainty to the percent-level would require precision measurements of the heavy higgsino-like neutralinos and charginos at TeV energies.

6 Acknowledgments

This work was supported in part by GDRI-ACPP of CNRS, the ANR project ToolsDMColl, the ‘SFB Transregio 33: The Dark Universe’, and grants RFBR-08-02-00856-a, RFBR-08-02-92499-a of the Russian Foundation for Basic Research.

References

- [1] Presentation:
<http://ilcagenda.linearcollider.org/contributionDisplay.py?contribId=196&sessionId=20&confId=2628>
- [2] WMAP, D. N. Spergel *et al.*, *Astrophys. J. Suppl.* **170**, 377 (2007);
SDSS, M. Tegmark *et al.*, *Phys. Rev.* **D69**, 103501 (2004);
- [3] Planck Collaboration, *astro-ph/0604069*.
- [4] M. M. Nojiri, G. Polesello and D. R. Tovey, *JHEP* **03**, 063 (2006).
- [5] E. A. Baltz, M. Battaglia, M. E. Peskin and T. Wizansky, *Phys. Rev.* **D74**, 103521 (2006).
- [6] T. Falk, K. A. Olive and M. Srednicki, *Phys. Lett.* **B354**, 99 (1995);
P. Gondolo and K. Freese, *JHEP* **07**, 052 (2002).
- [7] G. Bélanger, F. Boudjema, S. Kraml, A. Pukhov and A. Semenov, *Phys. Rev.* **D73**, 115007 (2006).
- [8] G. Bélanger, O. Kittel, S. Kraml, H. U. Martyn and A. Pukhov, *Phys. Rev. D* **78** (2008) 015011.
- [9] J. S. Lee *et al.*, *Comput. Phys. Commun.* **156**, 283 (2004).
- [10] G. Bélanger, F. Boudjema, A. Pukhov and A. Semenov, *Comput. Phys. Commun.* **176**, 367 (2007).
- [11] M. M. Nojiri, *Phys. Rev.* **D51**, 6281 (1995);
E. Boos *et al.*, *Eur. Phys. J.* **C30**, 395 (2003).
- [12] M. Pohl and H. J. Schreiber, *hep-ex/0206009*.
- [13] Y. Kizukuri and N. Oshimo, *Phys. Lett.* **B249**, 449 (1990);
A. Bartl, H. Fraas, O. Kittel and W. Majerotto, *Phys. Rev.* **D69**, 035007 (2004).